



Recent States and Prevention for Chronic Periodontitis and Diabetes in Actual Practice

Sakamoto D¹ and Bando H^{1,2,3,*}

¹Sakamoto Hospital, Higashi Kagawa city, Kagawa, Japan

²Tokushima University / Medical Research, Tokushima, Japan

³Japan Low Carbohydrate Diet Promotion Association, Kyoto, Japan

*Corresponding author: Bando H, Tokushima University / Medical Research, Nakashowa 1-61, Tokushima 770-0943, Japan; Tel: +81-90-3187-2485; E-mail: pianomed@bronze.ocn.ne.jp

Abstract

Chronic periodontitis and type 2 diabetes mellitus (T2DM) have been recently prevalent and crucial diseases. Both diseases show mutual vicious cycle as influencing each other. A systematic review revealed that T2DM cases with periodontitis have elevated odds ratio for retinopathy (2.8-8.7), neuropathy (3.2-6.6), nephropathy (1.9-8.5), cardiovascular complications (1.3-17.7) and mortality (2.3-8.5) compared to those without periodontitis. Adequate therapy for chronic periodontitis a few months would bring 0.27-0.48% decrease in HbA1c by meta-analyses. For patients with both diseases, low carbohydrate diet (LCD) seems to be effective associated with reduced HbA1c, body weight, Advanced Glycation End Products (AGEs) and inflammatory process.

Keywords: Chronic periodontitis; Type 2 diabetes mellitus (T2DM); Low carbohydrate diet (LCD); International Diabetes Federation (IDF); European Federation of Periodontology (EFP)

Commentary Article

In usual medical practice, type 2 diabetes mellitus (T2DM) and chronic periodontitis have been recently prevalent and crucial diseases worldwide [1]. These diseases are common in the departments of internal medicine and dentistry [2]. According to previous reports, both of these diseases may cause various complications as mutual vicious cycle [3]. A systematic review was conducted concerning the association of chronic periodontitis PD and diabetic complications [4]. It included 14 studies and analyzed diabetic macro- and microvascular complications and death with periodontitis. Compared to diabetes without periodontitis, higher risks were found in diabetic cases with periodontitis. The odds ratio revealed 2.8-8.7 in retinopathy, 3.2-6.6 in neuropathy, 1.9-8.5 in nephropathy, 1.3-17.7 in cardiovascular complications and 2.3-8.5 in mortality.

Regarding the problems of diabetes and periodontitis, dentists conducted clinical research of chairside screening for undiagnosed diabetes mellitus (UDM) for 7343 cases [5]. UDM was defined as no past diabetic history, or not fasting glucose >126 mg/dL, postprandial glucose > 200mg/dL. Periodontitis was

defined for positive medical history. As a result, the ratio of UCD was 5.6%, while obesity and edentulous (no teeth, toothless jaw) showed 12.6% and obesity and periodontitis showed 12.2%. In recent study, meta-analysis was reported concerning the association among nephropathy, retinopathy and periodontitis [1]. T2DM patients were included from 8 articles, in which 1207 cases with microvascular complications, and 1734 cases with periodontitis as well. As a result, odds ratio (OR) was 1.96 for periodontitis with microvascular complications. From subgroup analysis, the relationship existed among nephropathy, retinopathy and periodontitis for Asian race (OR 2.33) and North American populations (OR 1.42).

For current therapy of T2DM and chronic periodontitis, authors and colleagues have continued adequate medical practice with internal medicine and dentistry departments for years [2,6]. The guideline for T2DM is based on American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD), and others [7]. On the other hand, the guideline for T2DM and periodontitis is based on International Diabetes Federation (IDF) and European Federation of Periodontology (EFP) [8,9].

Received date: 01 June 2021; Accepted date: 06 June 2021; Published date: 12 June 2021

Citation: Sakamoto D, Bando H (2021). Recent States and Prevention for Chronic Periodontitis and Diabetes in Actual Practice. SunText Rev Dental Sci 2(2): 137.

DOI: <https://doi.org/10.51737/2766-4996.2021.037>

Copyright: © 2021 Sakamoto D, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

As to the basic therapy for diabetes, nutritional treatment would be important [10]. The standard method was formerly Calorie Restriction (CR), but recent tendency was changed. Low Carbohydrate Diet (LCD) was started in the medical regions by Drs. Atkins and Bernstein [11]. After that, LCD has been well-known in North American and European areas. For recent decade, clinical benefit of LCD has been prevalent to medical staffs and patients, which are also beneficial for glucose, lipid profile and arteriosclerotic cardiovascular disease (ASCVD) [12].

Our medical team have treated many patients with diabetes, obesity and chronic periodontitis [13,14]. Our group has initiated LCD at first in Japan, and developed LCD movement medically and socially through the activity of Japan LCD Promotion Association (JLCDPA) [15]. For practical use, three types of LCD have been recommended, which are super-LCD, standard-LCD, petite-LCD methods including carbohydrate 12%, 26%, 40%, respectively [16]. As any food includes a certain percentage of carbohydrate, and then minimum intake of carbohydrate becomes 12%. LCD has been applied to lots of cases with significant results. About 2900 cases showed 10% < weight reduction for 25%, and 5% < weight reduction for 59% [17]. Furthermore, LCD has brought hyperketonemia, which has been effective for not only protection of hyperglycemia, obesity, hyperlipidemia, but also reduction of Advanced Glycation End Products (AGEs) [18], inflammatory process and oxidation by reactive oxygen species (ROS) [19].

Among our clinical practice, a case with diabetes and chronic periapical periodontitis showed impressive course. He was 57-year-old man who suffered from impaired mastication and limited number of teeth [20]. By dental treatment and LCD for 3 months, general status was relieved satisfactory, in which weight decreased 100 kg to 90 kg and HbA1c improved 7.8% to 5.4%. Due to the standard guideline from EFP and IDF, the treatment for chronic periodontitis reveals clinical efficacy for diabetic patients [8]. By proper therapy for a few months, the reduction of HbA1c was reported to be about 0.27-0.48% by meta-analyses [21].

In the light of basic science, periodontal microbiome studies showed that the association between altered periodontal microbiome and changed glucose metabolism may be present [22]. Associated to this process, elevated glucose values will enhance the expression of pathogen receptors leading to stimulate response of the host. Hyperglycemia may also enhance pro-inflammatory response for the pathway of AGEs [22]. As periodontal pathogens product some cytokines for acute phase proteins, the insulin action and sensitivity may be impaired [23]. Discussion on the clinical efficacy has been continued concerning treatment of chronic periodontal disease only and/or combined therapy of periodontitis and diabetes. A recent study presented the comparison of HbA1c changes for patients with both diseases.

They were treated and followed by the treatment of full-mouth scaling associated with scaling root planing (SRP) for three months. The results showed that periodontal treatment had brought decreased inflammatory markers, indicating clinical efficacy of decreased HbA1c levels [23].

A recent study showed the correlation between glucose control and chronic periodontitis for T2DM patients [24]. The subjects included 182 cases with chronic periodontitis. As the protocol, several biomarkers were included such as fasting plasma glucose, HbA1c, remaining tooth number, probing depth (PD), bleeding on probing (BoP), clinical attachment level (CAL), and others. As a result, two groups (HbA1c is $\geq 7\%$ vs HbA1c $< 7\%$) revealed the significant difference as mean PD (3.78 vs. 3.42 mm, $p < 0.01$), mean CAL (4.5 vs. 4.1 mm, $p = 0.02$), mean remaining teeth (18.5 vs 20.4, $p < 0.01$). Consequently, extension of chronic periodontitis was found more prevalent in T2DM cases with poor glycemic variability than those with good control [24].

Conclusion

In conclusion, periodontal therapy can reduce HbA1c value. Diabetic patients have to check oral status and receive adequate periodontal treatment. Especially, when diabetic patient is hospitalized, to manage dental examination is required, because it is not included for the routine check [25]. Thus, physicians and medical staffs always have to keep dental evaluation in mind for diabetic patients.

Acknowledgments

None.

Funding

There was no funding received for this paper.

Conflicts of Interest

The authors declare no conflict of interest.

References

1. Wu HQ, Wei X, Yao JY, Qi JY, Xie HM, Sang AM, et al. Association between retinopathy, nephropathy, and periodontitis in type 2 diabetic patients: a Meta-analysis. *Int J Ophthalmol.* 2021; 14: 141-147.
2. Sakamoto D, Bando H. Recent trend from the clinical point of view for periodontitis and diabetes mellitus. *Dental Res Manag.* 2020; 4: 8-10.
3. Bando H. Diabetes Mellitus (DM) and Periodontal Disease (PD) with mutual vicious cycle. *Int J Res Stud Med Health Sci.* 2018; 3: 37-39.
4. Nguyen ATM, Akhter R, Garde S, Scott C, Twigg SM, Colagiuri S, et al. The association of periodontal disease with the

- complications of diabetes mellitus. A systematic review. *Diabetes Res Clin Pract.* 2020; 165: 108244.
5. Philips KH, Zhang S, Moss K, Ciarrocca K, Beck JD. Periodontal disease, undiagnosed diabetes, and body mass index: Implications for diabetes screening by dentists. *J Am Dent Assoc.* 2021; 152: 25-35.
 6. Bando H. Diabetes Mellitus (DM) and Periodontal Disease (PD) with Mutual Vicious Cycle. *Int J Res Stud Med Health Sci.* 2018; 3: 37-39.
 7. Buse JB, Wexle, DJ, Tsapas A, Rossing P, Mingrone G, Mathieu C, et al. 2019 Update to: Management of Hyperglycemia in Type 2 Diabetes, 2018. A Consensus Report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care.* 2019.
 8. Sanz M, Ceriello A, Buysschaert M. Scientific evidence on the links between periodontal diseases and diabetes: consensus report and guidelines of the joint EFP/IDF workshop on periodontal diseases and diabetes. *J Clin Periodontol.* 2018; 45: 138-149.
 9. Bissett SM, Presseau J, Rapley T, Preshaw PM. Uptake of best practice recommendations in the management of patients with diabetes and periodontitis: a cross-sectional survey of dental clinicians. *BDJ.* 2019; 226: 131-137.
 10. American Diabetes Association. 9. Pharmacologic approaches to glycemic treatment: Standards of Medical Care in Diabetes 2021. *Diabetes Care.* 2021; 44: 111-124
 11. Bernstein R. Dr. Bernstein's diabetes solution newly revised & updated. Little Brown and Company, New York, 2007.
 12. Tay J, Thompson CH, Luscombe-Marsh ND. Effects of an energy-restricted low carbohydrate, high unsaturated fat/low saturated fat diet versus a high-carbohydrate, low-fat diet in type 2 diabetes: A 2-year randomized clinical trial. *Diabetes Obes Metab.* 2018; 20: 858-871.
 13. Bando H, Ebe K, Muneta T, Bando M, Yonei Y. Difference of glucose variability between low carbohydrate diet (LCD) and calorie restriction (CR). *Asp Biomed Clin Case Rep.* 2018; 2: 4-15.
 14. Ebe K, Bando H, Muneta T, Bando M, Yonei Y. Remarkable improvement of glucose variability by Sodium-glucose cotransporter 2 (SGLT2) inhibitors using continuous glucose monitoring (CGM). *Diabetes Case Rep.* 2019; 4: 1.
 15. Muneta T, Kagaguchi E, Nagai Y, Matsumoto M, Ebe K, Watanabe H, et al. Ketone body elevation in placenta, umbilical cord, newborn and mother in normal delivery. *Glycative Stress Res.* 2016; 3: 133-140.
 16. Bando H. Useful tips for actual low carbohydrate diet (LCD) with super-, standard- and petit-lcd methods. *EC Nutrition.* 2020; 15: 01-04.
 17. Nakamura T, Bando H, Kawashima T, Dobashi M, Narita A. Weight reduction by effective protocol of diet and exercise. *J Diab Obes Metab.* 2019; 2:106.
 18. Perrone A, Giovino A, Benny J, Martinelli F. Advanced Glycation End Products (AGEs): biochemistry, signaling, analytical methods, and epigenetic effects. *Oxidative medicine and cellular longevity.* 2020; 2020: 1-18.
 19. Jakubczyk K, Dec K, Kałduńska J, Kawczuga D, Kochman J, Janda K. Reactive oxygen species - sources, functions, oxidative damage. *Pol Med J.* 2020; 48: 124-127.
 20. Sakamoto D, Bando H, Iwatsuki N, Sakamoto K, Okada M, Ogawa T. Remarkable improvement of periodontal diseases (PD) and diabetic state with weight reduction by super low carbohydrate diet (LCD). *J Oral Dent Health Res.* 2020; 2: 114.
 21. Madianos P, Koromantzos P. An update of the evidence on the potential impact of periodontal therapy on diabetes outcomes. *J Clin Periodontol.* 2018; 45: 188-195.
 22. Polak D, Sanui T, Nishimura F, Shapira L. Diabetes as a risk factor for periodontal disease-plausible mechanisms. *Periodontol.* 2020; 83: 46-58.
 23. Prasad R, Raj A, Khan SF, Agarwal S, Shalini B, Chandini HS. Effect of non-surgical periodontal therapy on glycemic control of patients with chronic periodontitis and type 2 diabetes mellitus: A 3 month follow-up intervention study. *Internat J Applied Dental Sci.* 2021; 7: 05-08.
 24. Stoicescu M, Calniceanu H, Țig I, Nemeth S, Tent A, Popa A, et al. Significant aspects and correlation between glycemic control and generalized chronic periodontitis in type 2 diabetes mellitus patients. *Experiment Therapeut Med.* 2021; 22: 671-675.
 25. Singh M, Bains VK, Jhingran R, Srivastava R, Madan R, Maurya SC, et al. Prevalence of periodontal disease in type 2 diabetes mellitus patients: A cross-sectional study. *Contemporary Clin Dentistry.* 2019; 10: 349.