



## Fundamental Sensory and Motor Neural Control in the Brain for the Musical Performance

Hiroshi BANDO<sup>1,2\*</sup>, Akiyo YOSHIOKA<sup>1</sup>, Yu NISHIKIORI<sup>1</sup>

<sup>1</sup>Shikoku Division of Integrative Medicine Japan (IMJ), Tokushima, Japan

<sup>2</sup>Tokushima University / Medical Research, Tokushima, Japan

Corresponding Author: **Hiroshi BANDO, MD, PhD, FACP** [ORCID ID](#)

**Address:** Tokushima University /Medical Research, Nakashowa 1-61, Tokushima 770-0943, Japan;

**Email:** [pianomed@bronze.ocn.ne.jp](mailto:pianomed@bronze.ocn.ne.jp)

**Received date:** 10 February 2022; **Accepted date:** 18 March 2022; **Published date:** 26 March 2022

**Citation:** Bando H, Yoshioka A, Nishikiori Y. Fundamental Sensory and Motor Neural Control in the Brain for the Musical Performance. J Health Care and Research. 2022 Mar 26;3(1):7-10.

**Copyright** © 2022 Bando H, Yoshioka A, Nishikiori Y. This is an open-access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium provided the original work is properly cited.

### Abstract

Music has beneficial power physically and psychologically. Among Integrative Medicine (IM), music therapy (MT) has been useful, and authors have continued research for IM, MT, and piano-playing. Most pianists do not consider the movement of their fingers, because the memorized process is transformed into automatic action. The function may involve the neural signals from the superior parietal lobule to the primary motor area and dorsal premotor cortex, which is called the sensory-motor transformations. The supplementary motor area (SMA) in the frontal lobe seems to be involved in the function of beat-based timing, expression, and activity of musical behavior.

### Keywords

Integrative Medicine, Music Therapy, Superior Parietal Lobule, Sensory-Motor Transformations, Supplementary Motor Area, Piano-Playing

### Abbreviations

IM: Integrative Medicine; MT: Music Therapy; SMA: Supplementary Motor Area

From a holistic point of view, Integrative Medicine (IM) has been developed worldwide [1]. Music has beneficial power to human beings physically and psychologically [2]. Authors and colleagues have managed various IM activities in the Shikoku Division of Integrative Medicine Japan (IMJ) and developed music therapy (MT) for years [3,4]. Several recent topics concerning MT, piano-playing, brain mechanism, and neuroimaging will be described in this article [5].

A comparative study using EEG was conducted among listening to neutral music, happy music, and control for 62 subjects [6]. As a result, happy music showed the efficacy of reducing anxiety and elevated functional connectivity of the right temporal lobe and occipital lobe. For brain research, the important condition requires three functions, correct sensory input, correct processing and determination, and output to the correct movement. Regarding this point, other cases with impaired dysfunction exist. For example, Parkinson's disease has involuntary

movements, and cerebral palsy has both contractions of flexor and extensor muscles simultaneously. If dystonia movement in the hand may be found for a pianist, it is difficult to play the piano smoothly [7,8].

When playing various music on the piano, most pianists do not consider the movement of their fingers, in which the movements are almost reflexive [9]. The reason would be that the memorized process is transformed into an automatic action [10]. The function may involve the neural signals from the superior parietal lobule to the primary motor area (M1) and dorsal premotor cortex, which is called the sensory-motor transformations [11] (Fig-1). Furthermore, this mechanism also contributes to the situation where a pianist can play a new music piece smoothly that is encountered first time [12].




Fig-1: Various area for sensory-motor transformations in the brain function

During playing the piano, each movement of the fingers is a single action of raising, spreading, and pushing the keyboard down. However, this performance is integrated into a huge and complex movement along the time axis using 10 fingers [13]. Continuing independent operations in the correct order is called "series operation". This is to arrange a variety of actions in the correct order along the time axis [14]. Such function has been related to a certain area in the cerebrum. Its responsive area would be mainly the primary motor area (M1) and supplementary motor area (SMA, supplementary

motor cortex) in the frontal lobe [15]. This area can contribute to the actual mechanism for complex and delicate movement for playing the piano. The function is thought to be stocking images of several actions together and executing the program in the correct order. SMA has been known as the crucial center for managing the planning of the program and performing the execution during playing the piano and listening to the music [16]. Music performance needs combined perspectives processing in visual, auditory, spatial, motor, and emotional domains. By integrating these multimodal factors, the SMA network may construct the internal expression of music performance.

As to the research of finger exercise, the changes in the excitability of M1 and SMA were investigated for 28 normal subjects [15]. The results showed the activity of increased M1 and decreased SMA. Consequently, it supposes the function of each of MA and SMA. In other words, rather strong exercise seems to give M1 protective efficacy and it gives opposite influence to SMA situated just adjacent to M1. SMA can work actively when starting new and complex movements, but on the other hand, it does not provide remarkable function when performing familiar movements one after another. Therefore, it is suggested that SMA contributes to the piano performance when the finger moves like never before in contact with a new music piece [16]. From the above, it is considered that SMA may have two functions. One is the start and control of complicated operations, and in particular, to build a program for what kind of operation is performed at adequate timing. The second is to switch the movement, which is considered to contribute to the creation of new complicated movements and procedures based on the memory conventionally already stored.

Several recent reports are found concerning the function of SMA. Regarding the research of neuroscientific field, investigations of musical creativity have brought the involved mechanisms of motor regions [17]. They include SMA, premotor cortex, posterior inferior frontal gyrus, which seem to be implicated for high level capacities concerning motor sequencing. As to regular rhythm in music, the presence of the background was studied for non-human primate. By maintaining the rhythm of visual

metronome, neural activity of SMA was recorded. SMA showed regular rhythmic gamma band bursts (30-40Hz) suggesting the existence of internal tempo center [18]. Consequently, SMA may have a role of dynamic metronome for rhythm and clock for total time. Regarding the brain mechanism composing creative music, comparative study was conducted between musicians and non-musicians for measuring evoked activation by functional magnetic resonance imaging (fMRI) [19]. The results showed that musicians revealed larger activation of the SMA, dorsolateral prefrontal cortex and anterior cingulate cortex. From these, SMA may play a role for the expression and activity of musical behavior. Music improvisation has been the complex creative behavior. Brain activity during improvisation was investigated using fMRI [20]. Compared with pre-learned certain melody for control piece, music improvisation showed stronger node activity in SMA, lateral premotor cortex, dorsolateral prefrontal cortex and Broca's area.

In summary, recent neuroimaging topics were introduced. Among them, SMA seems to be involved in the function of beat-based timing, expression and activity of musical behavior [21]. Research on music, MT, and piano performance from empirical, theoretical, and practical perspectives has been developed [22]. In the future, it will reduce people's mental stress and contribute to happiness of the people.

### Conflict of Interest

The authors have read and approved the final version of the manuscript. The authors have no conflicts of interest to declare.

### References

- [1] Ng JY, Jain A. Complementary and alternative medicine mention and recommendations in guidelines for anxiety: A systematic review and quality assessment. *Psychiatry Res*. 2022 Mar;309:114388. [PMID: 35051879]
- [2] Tobeta B, Bando H. Efficacy of sound treatment for better meditation and sleep in music therapy. *Int J Complement Alt Med*. 2021;14(4):192-94.
- [3] Bando H (2021). Cross Modality Leading to Sonic Seasoning with Music for Integrative Medicine (IM).

*Int J Conf Proc*. 2021;2(5):ICP.000547.

- [4] Yoshioka A, Nishikiori Y, Bando H (2020) Music Therapy Session with Various Elements for Clinical Effects and Comfortable Mood. *Biomed Sci J*. 2020;2:11.
- [5] Yamashita M, Ohsawa C, Suzuki M, Guo X, Sadakata M, Otsuka Y, Asano K, Abe N, Sekiyama K. Neural Advantages of Older Musicians Involve the Cerebellum: Implications for Healthy Aging Through Lifelong Musical Instrument Training. *Front Hum Neurosci*. 2022 Jan 5;15:784026. [PMID: 35069154]
- [6] Huang B, Hao X, Long S, Ding R, Wang J, Liu Y, Guo S, Lu J, He M, Yao D. The Benefits of Music Listening for Induced State Anxiety: Behavioral and Physiological Evidence. *Brain Sci*. 2021 Oct 9;11(10):1332. [PMID: 34679397]
- [7] Kita K, Furuya S, Osu R, Sakamoto T, Hanakawa T. Aberrant Cerebello-Cortical Connectivity in Pianists With Focal Task-Specific Dystonia. *Cereb Cortex*. 2021 Aug 26;31(10):4853-63. [PMID: 34013319]
- [8] Lee SH, Sanchez-Ramos J, Murtagh R, Vu T, Hardwick D, Carey SL. A Case Report: Using Multimodalities to Examine a Professional Pianist with Focal Dystonia. In *Perspectives in Performing Arts Medicine Practice 2020* (pp. 165-178). Springer, Cham.
- [9] Olszewska AM, Gaca M, Herman AM, Jednoróg K, Marchewka A. How Musical Training Shapes the Adult Brain: Predispositions and Neuroplasticity. *Front Neurosci*. 2021 Mar 10;15:630829. [PMID: 33776638]
- [10] Iorio C, Brattico E, Munk Larsen F, Vuust P, Bonetti L. The effect of mental practice on music memorization. *Psychology of Music*. 2022 Jan;50(1):230-44.
- [11] Zareian B, Zhang Z, Zagha E. Cortical Localization of the Sensory-Motor Transformation in a Whisker Detection Task in Mice. *eNeuro*. 2021 Feb 1;8(1):ENEURO.0004-21.2021. [PMID: 33495240]
- [12] Fasano MC, Gleran E, Gold BP, Sheng D, Sams M, Vuust P, Rauschecker JP, Brattico E. Inter-subject Similarity of Brain Activity in Expert Musicians After Multimodal Learning: A Behavioral and Neuroimaging Study on Learning to Play a Piano Sonata. *Neuroscience*. 2020 Aug 10;441:102-16. [PMID: 32569807]
- [13] Tokgoz S, Aydogdu D, Ilhan B, Sahin Y, Bariseri N, Ozturkler BM, Çukur T. Musical mirror-symmetrical movement tasks: comparison of rhythm versus melody-playing. *Neuroreport*. 2020 May 7;31(7):523-

29. [PMID: 32221114]
- [14] Yang Y. Piano performance and music automatic notation algorithm teaching system based on artificial intelligence. *Mobile Information Systems*. 2021 Oct 19;2021.
- [15] Coco M, Perciavalle V, Cavallari P, Perciavalle V. Effects of an Exhaustive Exercise on Motor Skill Learning and on the Excitability of Primary Motor Cortex and Supplementary Motor Area. *Medicine (Baltimore)*. 2016 Mar;95(11):e2978. [PMID: 26986109]
- [16] Tanaka S, Kirino E. Dynamic Reconfiguration of the Supplementary Motor Area Network during Imagined Music Performance. *Front Hum Neurosci*. 2017 Dec 12;11:606. [PMID: 29311870]
- [17] Bashwiner D, Bacon D. Musical creativity and the motor system. *Current Opinion in Behavioral Sciences*. 2019 Jun 1;27:146-53.
- [18] Cadena-Valencia J, García-Garibay O, Merchant H, Jazayeri M, de Lafuente V. Entrainment and maintenance of an internal metronome in supplementary motor area. *Elife*. 2018 Oct 22;7:e38983. [PMID: 30346275]
- [19] de Aquino MPB, Verdejo-Román J, Pérez-García M, Pérez-García P. Different role of the supplementary motor area and the insula between musicians and non-musicians in a controlled musical creativity task. *Sci Rep*. 2019 Sep 10;9(1):13006. [PMID: 31506553]
- [20] Dhakal K, Norgaard M, Adhikari BM, Yun KS, Dhamala M. Higher Node Activity with Less Functional Connectivity During Musical Improvisation. *Brain Connect*. 2019 Apr;9(3):296-309. [PMID: 30618291]
- [21] Leow LA, Rinchon C, Emerick M, Grahn J. Supplementary motor area contributions to rhythm perception. *bioRxiv*. 2021 Jan 1.
- [22] de Witte M, Knapen A, Stams GJ, Moonen X, van Hooren S. Development of a music therapy micro-intervention for stress reduction. *The Arts in Psychotherapy*. 2022 Feb 1;77:101872.

